



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DEATHS DURING WEEK ENDED FEBRUARY 15, 1919, IN CITIES.

The following table shows the registered deaths from all causes and from pneumonia (all forms) and influenza combined, in certain large cities of the United States during the week ended February 15, 1919. The annual death rates per 1,000 population for the week and for the corresponding week of previous years are also shown.

The data are taken from the "Weekly Health Index," February 18, 1919, issued by the Bureau of the Census, Department of Commerce. The populations used in computing the rates are estimated by the Bureau of the Census as of July 1, 1918.

Registered deaths and annual death rates per 1,000 population in certain large cities of the United States, week ended February 15, 1919—Deaths from all causes, and from pneumonia (all forms) and influenza combined.

City.	Population July 1, 1918, estimated.	Total deaths, all causes.	Annual death rate per 1,000.	Annual death rate for preceding years. ¹	Influenza and pneumonia (all forms).	
					Number of deaths.	Annual rate per 1,000.
Albany, N. Y.	112,565	47	21.8	C 19.0	16	7.4
Atlanta, Ga.	201,732	72	18.6	C 19.1		
Baltimore, Md.	669,981	313	24.4	A 20.0	117	9.1
Birmingham, Ala.	197,670	84	22.2	A 18.8	28	7.4
Boston, Mass.	785,245	262	17.4	A 18.0	71	4.7
Buffalo, N. Y.	473,229	187	20.6	C 15.3	35	3.9
Cambridge, Mass.	111,433	28	13.1	A 16.6	3	1.4
Chicago, Ill.	2,596,681	903	18.1	A 17.3	235	4.7
Cincinnati, Ohio.	418,022	162	20.2	C 21.2	37	4.6
Cleveland, Ohio.	810,306	249	16.0	C 12.2	80	5.2
Columbus, Ohio.	225,296	68	15.7	C 21.3	15	3.5
Dayton, Ohio.	130,655	38	15.2	C 15.6		
Denver, Colo.		101				
Fall River, Mass.	128,392	41	16.7	C 18.7	15	6.1
Grand Rapids, Mich.	135,450	33	12.7	C 9.6		
Indianapolis, Ind.	289,577	98	17.6	C 13.9		
Kansas City, Mo.	313,785	134	22.3	C 20.6	51	8.5
Los Angeles, Cal.	568,495	160	14.7	A 15.3	21	1.9
Louisville, Ky.	242,707	85	18.3	C 21.7	19	4.1
Lowell, Mass.	109,081	38	18.2	A 19.3	18	8.6
Memphis, Tenn.	154,759	49	16.5	C 15.5	19	6.4
Milwaukee, Wis.	453,481	110	12.6	A 14.7		
Minneapolis, Minn.	383,442	95	12.9	C 14.1		
Nashville, Tenn.	119,215	59	25.8	C 21.0	16	7.0
Newark, N. J.	428,684	132	16.1	C 17.8	32	3.9
New Haven, Conn.	154,865	52	17.5	C 20.9	11	3.7
New Orleans, La.	382,273	176	24.0	A 22.4	49	6.7
New York, N. Y.	5,215,879	2,090	20.9	C 17.4	786	7.9
Omaha, Neb.	180,264	47	13.6	C 8.1		
Philadelphia, Pa.	1,761,371	728	21.6	18.5	232	6.9
Pittsburgh, Pa.	593,303	309	27.2	C 18.7	163	14.3
Portland, Oreg.		64			10	
Providence, R. I.	263,613	103	20.4	C 19.0	28	5.5
Richmond, Va.	160,719	69	22.4	C 21.7	9	2.9
Rochester, N. Y.	264,856	68	13.4	C 15.4	16	3.2
St. Louis, Mo.	779,951	239	16.0	C 18.7	64	4.3
St. Paul, Minn.	257,699	58	11.7	C 7.1		
San Francisco, Cal.	478,530	151	16.5	C 18.1	20	2.2
Seattle, Wash.		79			21	
Spokane, Wash.		22				
Syracuse, N. Y.	161,404	57	18.4	C 20.4	10	3.2
Toledo, Ohio.	262,234	70	13.9	A 13.6	11	2.2
Washington, D. C.	401,681	149	19.3	A 19.1	40	5.2
Worcester, Mass.	173,650	87	26.1	C 19.8	23	6.9

¹ "A" indicates that the rate given is the average annual death rate per 1,000 population for the corresponding week of the years 1913 to 1917, inclusive. "C" indicates that the rate is the annual death rate per 1,000 population for the corresponding week of 1918.

² Population estimated as of July 1, 1919.

³ Rate is based on statistics of 1915, 1916, and 1917.